

MANIFOLDS COVERED BY LINES AND EXTREMAL RAYS

CARLA NOVELLI AND GIANLUCA OCCHETTA

ABSTRACT. Let X be a smooth complex projective variety and let $H \in \text{Pic}(X)$ be an ample line bundle. Assume that X is covered by rational curves with degree one with respect to H and with anticanonical degree greater than or equal to $(\dim X - 1)/2$. We prove that there is a covering family of such curves whose numerical class spans an extremal ray in the cone of curves $\text{NE}(X)$.

INTRODUCTION

Let X be a smooth complex projective variety which admits a morphism $\varphi: X \rightarrow Z$ onto a normal variety Z such that the anticanonical bundle $-K_X$ is φ -ample, $\dim X > \dim Z$ and $\rho_X = \rho_Z + 1$ (*i.e.* an elementary extremal contraction of fiber type).

It is well known, by fundamental results of Mori theory, that through every point of X there is a rational curve contracted by φ . The numerical classes of these curves lie in an extremal ray of the cone $\text{NE}(X)$. By taking a covering family of such curves, with minimal degree with respect to some fixed ample line bundle, one obtains a *quasi-unsplit* family of rational curves, *i.e.* a family such that the reducible components of all the degenerations of curves in the family are numerically proportional to a curve in the family. It is very natural to ask if the converse is also true:

Given a covering quasi-unsplit family V of rational curves, is there an extremal elementary contraction which contracts all curves in the family or, in other words, does the numerical class of a general curve in the family span an extremal ray of $\text{NE}(X)$?

By [8] (see also [14] and [10]) there is always a rational fibration, defined on an open set of X , whose general fibers are proper, which contracts a general curve in V . More precisely, a general fiber is an equivalence class with respect to the relation induced by the closure $\mathcal{V}$ of the family V in the Chow scheme of X in the following way: two points x and y are equivalent if there exists a connected chain of cycles in $\mathcal{V}$ which joins x and y .

By a careful study of this fibration and of its indeterminacy locus, a partial answer to this question has been given in [6, Theorem 2]; namely, if the dimension of a general equivalence class is greater than or equal to the dimension of the variety minus three then the numerical class of a general curve in the family spans an extremal ray of $\text{NE}(X)$.

Before the results in [6] a special but very natural situation in which the question arises has been studied in [5]. In that paper manifolds covered by rational curves

2000 *Mathematics Subject Classification.* 14J40, 14E30, 14C99.

Key words and phrases. Rational curves, extremal rays.

of degree one with respect to an ample line bundle H were considered, and it was proved that a covering family of such curves (we will call them lines, by abuse) of anticanonical degree greater than or equal to $\frac{\dim X + 3}{2}$ spans an extremal ray (see also [4, Theorem 2.4]).

Recently, in [15, Theorem 7.3], the extremality of a covering family V of lines was proved under the weaker assumption that the anticanonical degree of such curves, which we will denote by abuse of notation by $-K_X \cdot V$, is greater than or equal to $\frac{\dim X + 1}{2}$.

The goal of the present paper is to prove the following

Theorem. *Let (X, H) be a polarized manifold with a dominating family of rational curves V such that $H \cdot V = 1$. If $-K_X \cdot V \geq \frac{\dim X - 1}{2}$, then $[V]$ spans an extremal ray of $\text{NE}(X)$.*

The main idea is, as in [15], to combine the ideas and techniques of [5], especially taking into consideration a suitable adjoint divisor $K_X + mH$ and studying its nefness, with those of [6], in particular regarding the existence of special curves in the indeterminacy locus of the rational fibration associated to V .

1. BACKGROUND MATERIAL

Let X be a smooth projective variety defined over the field of complex numbers. A contraction $\varphi: X \rightarrow Z$ is a proper surjective map with connected fibers onto a normal variety Z .

If the canonical bundle K_X is not nef, then the negative part of the cone $\text{NE}(X)$ of effective 1-cycles is locally polyhedral, by the Cone Theorem. By the Contraction Theorem, to every face in this part of the cone is associated a contraction, called *extremal contraction* or *Fano–Mori contraction*.

An extremal contraction associated to a face of dimension one, *i.e.* to an extremal ray, is called an *elementary contraction*; an extremal ray R is called **numerically effective**, and the associated contraction is said to be of **fiber type**, if $\dim Z < \dim X$; otherwise the ray is called **non nef** and the contraction is **birational**.

If the codimension of the exceptional locus of an elementary birational contraction is equal to one, the ray and the contraction are called **divisorial**, otherwise they are called **small**.

A Cartier divisor which is the pull-back of an ample divisor A on Z is called a *supporting divisor* of the contraction φ .

If the anticanonical bundle of X is ample, X is called a Fano manifold. For a Fano manifold, the *index*, denoted by r_X , is defined as the largest natural number r such that $-K_X = rH$ for some (ample) divisor H on X .

Definition 1.1. A *family of rational curves* is an irreducible component $V \subset \text{Ratcurves}^n(X)$ (see [14, Definition 2.11]). Given a rational curve we will call a *family of deformations* of that curve any irreducible component of $\text{Ratcurves}^n(X)$ containing the point parametrizing that curve.

We will say that V is *unsplit* if it is proper.

We define $\text{Locus}(V)$ to be the set of points of X through which there is a curve among those parametrized by V ; we say that V is a *covering family* if $\text{Locus}(V) = X$ and that V is a *dominating family* if $\overline{\text{Locus}(V)} = X$.

We denote by V_x the subscheme of V parametrizing rational curves passing through

$x \in \text{Locus}(V)$ and by $\text{Locus}(V_x)$ the set of points of X through which there is a curve among those parametrized by V_x .

By abuse of notation, given a line bundle $L \in \text{Pic}(X)$, we will denote by $L \cdot V$ the intersection number $L \cdot C_V$, with C_V any curve among those parametrized by V .

Proposition 1.2. ([14, IV.2.6]) *Let V be an unsplit family of rational curves on X . Then*

- (a) $\dim X - K_X \cdot V \leq \dim \text{Locus}(V) + \dim \text{Locus}(V_x) + 1$;
- (b) $-K_X \cdot V \leq \dim \text{Locus}(V_x) + 1$.

This last proposition, in case V is the unsplit family of deformations of a rational curve of minimal anticanonical degree in an extremal face of $\text{NE}(X)$, gives the *fiber locus inequality*.

Proposition 1.3. ([12, Theorem 0.4], [18, Theorem 1.1]) *Let φ be a Fano–Mori contraction of X . Denote by E the exceptional locus of φ and by F an irreducible component of a non-trivial fiber of φ . Then*

$$\dim E + \dim F \geq \dim X + l - 1,$$

where $l := \min\{-K_X \cdot C \mid C \text{ is a rational curve in } F\}$. If φ is the contraction of an extremal ray R , then $l(R) := l$ is called the length of the ray.

Definition 1.4. We define a *Chow family of rational curves* $\mathcal{W}$ to be an irreducible component of $\text{Chow}(X)$ parametrizing rational and connected 1-cycles.

We define $\text{Locus}(\mathcal{W})$ to be the set of points of X through which there is a cycle among those parametrized by $\mathcal{W}$; notice that $\text{Locus}(\mathcal{W})$ is a closed subset of X ([14, II.2.3]). We say that $\mathcal{W}$ is a *covering family* if $\text{Locus}(\mathcal{W}) = X$.

Definition 1.5. If V is a family of rational curves, the closure of the image of V in $\text{Chow}(X)$, denoted by $\mathcal{V}$, is called the *Chow family associated to V* .

Remark 1.6. If V is proper, i.e. if the family is unsplit, then V corresponds to the normalization of the associated Chow family $\mathcal{V}$.

Definition 1.7. Let $\mathcal{V}$ be the Chow family associated to a family of rational curves V . We say that V is *quasi-unsplit* if every component of any reducible cycle in $\mathcal{V}$ is numerically proportional to V .

Definition 1.8. Let $\mathcal{W}$ be a Chow family of rational curves on X and $Z \subset X$. We define $\text{Locus}(\mathcal{W})_Z$ to be the set of points $x \in X$ such that there exists a cycle Γ among those parametrized by $\mathcal{W}$ with $\Gamma \cap Z \neq \emptyset$ and $x \in \Gamma$.

We define $\text{ChLocus}(\mathcal{W})_Z$ to be the set of points $x \in X$ such that there exists a chain of cycles among those parametrized by $\mathcal{W}$ connecting x and Z .

Lemma 1.9. ([14, Proposition IV.3.13.3], [1, Lemma 4.1]) *Let $Z \subset X$ be a closed subset and let $\mathcal{W}$ be a Chow family of rational curves. Then every curve contained in $\text{ChLocus}(\mathcal{W})_Z$ is numerically equivalent to a linear combination with rational coefficients of a curve contained in Z and irreducible components of cycles among those parametrized by $\mathcal{W}$ which intersect Z .*

In particular, if V is a quasi-unsplit family, every curve contained in $\text{ChLocus}(\mathcal{V})_Z$ is numerically equivalent to a linear combination with rational coefficients of a curve contained in Z and a curve among those parametrized by V . We will write

$$N_1^X(\text{ChLocus}(\mathcal{V})_Z) = \langle \text{NE}(Z), [V] \rangle.$$

Lemma 1.10. (Cf. [5, Proof of Lemma 1.4.5], [16, Lemma 1]) *Let $Z \subset X$ be a closed subset and let V be a quasi-unsplit family of rational curves. Then every curve contained in $\text{ChLocus}(\mathcal{V})_Z$ is numerically equivalent to a linear combination with rational coefficients*

$$\lambda C_Z + \mu C_V,$$

where C_Z is a curve in Z , C_V is a curve among those parametrized by V and $\lambda \geq 0$.

Corollary 1.11. (Cf. [9, Corollary 2.2 and Remark 2.4]) *Let Σ be an extremal face of $\text{NE}(X)$ and denote by F a fiber of the contraction associated to Σ . Let V be a quasi-unsplit family numerically independent from curves whose numerical class is in Σ . Then*

$$\text{NE}^X(\text{ChLocus}(\mathcal{V})_F) = \langle \Sigma, [V] \rangle,$$

i.e. the numerical class in X of a curve in $\text{ChLocus}(\mathcal{V})_F$ is in the subcone of $\text{NE}(X)$ generated by Σ and $[V]$.

Lemma 1.12. *Let D an effective divisor on X and L a nef divisor. If $(L + D)|_D$ is nef then $L + D$ is nef.*

Proof. Assume that γ is an effective curve on X such that $(L + D) \cdot \gamma < 0$. By the nefness of L we have $D \cdot \gamma < 0$, hence $\gamma \subset D$. But $L + D$ is nef on D , a contradiction. $\square$

2. RATIONALLY CONNECTED FIBRATIONS

Let X be a smooth complex projective variety and let $\mathcal{W}$ be a covering Chow family of rational curves.

Definition 2.1. The family $\mathcal{W}$ defines a relation of rational connectedness with respect to $\mathcal{W}$, which we shall call *rc($\mathcal{W}$)-relation* for short, in the following way: x and y are in *rc($\mathcal{W}$)-relation* if there exists a chain of cycles among those parametrized by $\mathcal{W}$ which joins x and y .

To the *rc($\mathcal{W}$)-relation* we can associate a fibration, at least on an open subset ([7], [14, IV.4.16]); we will call it *rc($\mathcal{W}$)-fibration*.

We have a diagram, coming from the universal family over $\text{Chow}(X)$:

$$\begin{array}{ccc} \mathcal{U} & \xrightarrow{i} & X \\ p \downarrow & & \\ \mathcal{W}. & & \end{array}$$

In the above diagram, the map i is induced by the evaluation and the fibers of p are connected with rational components. Moreover, both i and p are proper (see for instance [14, II.2.2]).

In the notation of [6], by [10, Theorem 5.9] there exists a closed irreducible subset of $\text{Chow}(X)$ such that, denoting by Y its normalization and by $Z \subset Y \times X$ the restriction of the universal family, we have a commutative diagram

$$(2.1.1) \quad \begin{array}{ccc} Z & \xrightarrow{e} & X \\ p \downarrow & \swarrow q & \\ Y & & \end{array}$$

where e is a birational morphism whose exceptional locus E does not dominate Y . Moreover, a general fiber of q is irreducible and is a $\text{rc}(\mathcal{W})$ -equivalence class. Let B be the image of E in X ; note that $\dim B \leq \dim X - 2$, as X is smooth.

If we consider a (covering) Chow family $\mathcal{V}$, associated to a quasi-unsplit dominating family V , then by [6, Proposition 1, (ii)] B is the union of all $\text{rc}(\mathcal{V})$ -equivalence classes of dimension greater than $\dim X - \dim Y$.

Moreover we have the following

Lemma 2.2. *Let V be a quasi-unsplit dominating family of rational curves on a smooth complex projective variety X . Let B be the indeterminacy locus of the $\text{rc}(\mathcal{V})$ -fibration $q: X \dashrightarrow Y$, let D be a very ample divisor on $q(X \setminus B)$ and let $\widehat{D} := q^{-1}D$. Then*

- (1) $\widehat{D} \cdot V = 0$;
- (2) if $C \not\subset B$ is a curve whose numerical class is not proportional to $[V]$, then $\widehat{D} \cdot C > 0$;
- (3) if $\widehat{D} \cdot C > 0$ for every curve $C \subset B$ whose class is not proportional to $[V]$, then $[V]$ spans an extremal ray of $\text{NE}(X)$.

Proof. A general cycle among those parametrized by $\mathcal{V}$ is contained in a fiber of q disjoint from $\widehat{D}$, so $\widehat{D} \cdot V = 0$.

If C is as in (2), then $q(C)$ is a curve in Y and the result follows from the projection formula.

Finally, if $[V]$ does not span an extremal ray, then either $\widehat{D}$ is not nef or $\widehat{D}$ is nef but $\widehat{D}^\perp \cap \text{NE}(X) \supsetneq [V]$. In both cases there exists a curve $C \subset X$ whose class is not proportional to $[V]$ such that $\widehat{D} \cdot C \leq 0$. Such a curve must be contained in B by [6, Proof of Proposition 1]. $\square$

Corollary 2.3. *Let V be a quasi-unsplit dominating family of rational curves on a smooth complex projective variety X ; denote by B the indeterminacy locus of the $\text{rc}(\mathcal{V})$ -fibration and by f_V the dimension of the general $\text{rc}(\mathcal{V})$ -equivalence class. If $[V]$ does not span an extremal ray of $\text{NE}(X)$, then B is not empty. In particular there exist $\text{rc}(\mathcal{V})$ -equivalence classes of dimension $\geq f_V + 1$.*

We now give a lower bound on the dimension of $\text{ChLocus}(\mathcal{V})_S$, depending on the position the subvariety S with respect to the indeterminacy locus of the $\text{rc}(\mathcal{V})$ -fibration.

Lemma 2.4. *Let V be a quasi-unsplit dominating family of rational curves on a smooth complex projective variety X ; denote by B the indeterminacy locus of the $\text{rc}(\mathcal{V})$ -fibration and by f_V the dimension of the general $\text{rc}(\mathcal{V})$ -equivalence class. Let $S \subset X$ be an irreducible subvariety such that $[V] \notin \text{NE}^X(S)$. Then there exists an irreducible component X_S of $\text{ChLocus}(\mathcal{V})_S$ containing S and such that*

- (1) if $S \not\subset B$, then $\dim X_S \geq \dim S + f_V$.
- (2) if $S \subset B$, then $\dim X_S \geq \dim S + f_V + 1$.

Proof. We refer to diagram (2.1.1). Let $S' \subset Z$ be an irreducible component of $e^{-1}(S)$ which dominates S via e . By our assumptions S' meets any fiber of p at most in points, hence the subset $Z_S := p^{-1}(p(S'))$ has dimension

$$(2.4.1) \quad \dim Z_S = \dim p(S') + f_V = \dim S' + f_V \geq \dim S + f_V.$$

Moreover, notice that $S \subset e(Z_S) \subset \text{ChLocus}(\mathcal{V})_S$.

Assume that $S \not\subset B$. Then $Z_S \not\subset E$, hence the map $e|_{Z_S}: Z_S \rightarrow X$ is generically finite. Therefore, in view of (2.4.1), $\dim e(Z_S) = \dim Z_S \geq \dim S + f_V$, and so (1) is proved.

Assume now that $S \subset B$. As $e(Z_S) \subset \text{ChLocus}(\mathcal{V})_S$ and B is closed with respect to $\text{rc}(\mathcal{V})$ -equivalence, we have $e(Z_S) \subset B$.

Assertion (2) will follow once we prove that the general fiber G of $e|_{Z_S}$ has dimension strictly smaller than the general fiber of $e|_{S'}$. In fact, recalling also (2.4.1), in this case we will have

$$\dim e(Z_S) = \dim Z_S - \dim G > (\dim S' + f_V) - (\dim S' - \dim S) = f_V + \dim S.$$

Let $S^1 \subset S'$ be the closed subset on which e is not equidimensional, let $Z_{S^1} := p^{-1}(p(S^1))$, and let x be a general point in $e(Z_S) \setminus e(Z_{S^1})$. Let G be the fiber of $e|_{Z_S}$ over x and let Z_G be $p^{-1}(p(G))$. Since $e(Z_G) \subset \text{ChLocus}(\mathcal{V})_x$, the image via e of any curve in $Z_G \cap S'$ must be a point, otherwise it would be a curve in $S \cap \text{ChLocus}(\mathcal{V})_x$, which is a contradiction, since curves in S are numerically independent from $[V]$.

As a consequence, $Z_G \cap S'$ is contained in a finite union of fibers $F_1, \dots, F_k$ of e over points $s_1, \dots, s_k$. By our choice of x the dimension of these fibers is $\dim S' - \dim S$. Let y be a general point of Z_G and denote by y' a point in $p^{-1}(p(y)) \cap S'$; y' is contained in a fiber of e over one of the points $s_1, \dots, s_k$. The fiber G_y of $e|_{Z_S}$ containing y is thus contained in $p^{-1}(p(F_i))$ for some $i = 1, \dots, k$.

It follows that the dimension of the general fiber of $e|_{Z_S}$ is the dimension of the general fiber of $e|_{Z_F}$, where $Z_F := p^{-1}(p(\cup F_i))$. Notice that $\dim e(Z_F) > f_V$, otherwise every fiber of p contained in Z_F would have the same image via e .

Therefore

$$\dim G = \dim Z_F - \dim e(Z_F) < (\dim F + f_V) - f_V = \dim F = \dim S' - \dim S,$$

and we are done. $\square$

Remark 2.5. Both the bounds in Lemma (2.4) are sharp. An example for the second one is given by [6, Example 2]: in that example $B \simeq \mathbb{P}^2 \times \mathbb{P}^1$; taking as S a fiber of the projection onto $\mathbb{P}^2$ we have equality in (2).

3. BLOWING-DOWN

In this section we consider the following situation, which will show up in the proof of Theorem (4.3):

Lemma 3.1. *Let (X, H) be a polarized manifold with a dominating family of rational curves V such that $H \cdot V = 1$. Denote by f_V the dimension of the general $\text{rc}(\mathcal{V})$ -equivalence class and assume that there exists an extremal face Σ in $\text{NE}(X)$ whose associate contraction $\sigma: X \rightarrow X'$ is a smooth blow-up along a disjoint union of subvarieties T_i of dimension $\leq f_V$ such that $E_i \cdot V = 0$ for every exceptional divisor E_i .*

Finally denote by V' a family of deformation of $\sigma(C)$, with C a general curve parametrized by V . Then

- (1) $-K_{X'} \cdot V' = -K_X \cdot V$;
- (2) *there exists an ample line bundle H' on X' such that $H' \cdot V' = 1$;*
- (3) *if C' is a curve parametrized by V' such that $T_i \cap C' \neq \emptyset$, then $C' \subset T_i$;*
- (4) $\rho_{X'} > 1$;

- (5) if $[V']$ spans an extremal ray of $\text{NE}(X')$, then $[V]$ spans an extremal ray of $\text{NE}(X)$.

Proof. It is enough to prove the statement in case $\dim \Sigma = 1$, i.e. $\sigma: X \rightarrow X'$ is the blow-up of X' along a smooth subvariety T associated to the extremal ray Σ . In fact, if $\dim \Sigma > 1$, the contraction of Σ factors through elementary contractions, each one satisfying the assumptions in the statement.

Denote by E the exceptional locus of σ . Since $E \cdot V = 0$ the first assertion in the statement follows from the canonical bundle formula for blow-ups.

Moreover, the fact that $E \cdot V = 0$ also implies that any $\text{rc}(\mathcal{V})$ -equivalence class meeting E is actually contained in E . Therefore, if F is a non-trivial fiber of σ , then $\text{ChLocus}(\mathcal{V})_F \subseteq E$. By Lemma (2.4)

$$\dim \text{ChLocus}(\mathcal{V})_F \geq f_V + \dim F \geq \dim X - 1,$$

hence $E = \text{ChLocus}(\mathcal{V})_F$ and $\dim T = f_V$. So $\text{NE}^X(E) = \langle [V], \Sigma \rangle$ by Corollary (1.11).

The line bundle $(H + E)|_E$ is nef and it is trivial only on Σ , since $(H + E) \cdot \Sigma = 0$ and $(H + E) \cdot V = 1$. Then $H + E$ is nef by Lemma (1.12).

Notice also that $H + E$ is trivial only on Σ . Indeed, let γ be an effective curve on X such that $(H + E) \cdot \gamma = 0$. Due to the ampleness of H we have $E \cdot \gamma < 0$, hence $\gamma \subset E$. This implies that $[\gamma] \in \Sigma$.

Therefore $H + E = \sigma^* H'$, with H' an ample line bundle on X' . By the projection formula $H' \cdot V' = 1$, hence part (2) in the statement is proved.

Now, let C' be a curve of V' meeting T and assume by contradiction that C' is not contained in T ; denote by $\tilde{C}'$ its strict transform. Then

$$1 = H' \cdot C' = \sigma^* H' \cdot \tilde{C}' = (H + E) \cdot \tilde{C}' \geq 2,$$

which is a contradiction. It follows that every curve parametrized by V' which meets T is contained in it; so we get part (3) in the statement.

As to part (4), assume by contradiction that $\rho_{X'} = 1$. This implies that X' is $\text{rc}(\mathcal{V})$ -connected, but this is impossible as, in view of part (3), we cannot join points of T and points outside of T with curves parametrized by V' .

Assume now that $[V']$ spans an extremal ray of X' . Let B be the indeterminacy locus of the $\text{rc}(\mathcal{V})$ -fibration.

We claim that $E \cap B = \emptyset$.

Assume by contradiction that this is not the case; then E meets (and hence contains) an $\text{rc}(\mathcal{V})$ -equivalence class G of dimension $\dim G \geq f_V + 1$. Take a fiber F of σ meeting G . Then $\dim F + \dim G > \dim E$. On the other hand, $\dim(F \cap G) = 0$ as $[V] \notin \Sigma$. So we get a contradiction.

Let A be a supporting divisor of the contraction associated to $[V']$. The pull-back $\sigma^* A$ defines a two-dimensional face Π of $\text{NE}(X)$ containing Σ and $[V]$. Assume that Π is not spanned by Σ and $[V]$; in this case there exists a curve C whose class is in Π and such that $E \cdot C > 0$. Let $\hat{D}$ be as in Lemma (2.2); by the same lemma $\hat{D} \cdot \Sigma > 0$ and $\hat{D} \cdot V = 0$, hence we have $\hat{D} \cdot C < 0$ and so $C \subset B$, contradicting $E \cap B = \emptyset$. $\square$

4. MAIN THEOREM

First of all we consider polarized manifolds (X, H) with a quasi-unsplit dominating family of rational curves V proving that if, for m large enough, the adjoint divisor $K_X + mH$ defines an extremal face containing $[V]$, then $[V]$ spans an extremal ray of X .

Proposition 4.1. *Let (X, H) be a polarized manifold which admits a quasi-unsplit dominating family of rational curves V ; denote by f_V the dimension of a general $\text{rc}(\mathcal{V})$ -equivalence class.*

If, for some integer m such that $m + f_V \geq \dim X - 3$, the divisor $K_X + mH$ is nef and it is trivial on $[V]$, then $[V]$ spans an extremal ray of $\text{NE}(X)$.

Proof. Assume by contradiction that $[V]$ does not span an extremal ray in $\text{NE}(X)$. This implies that $K_X + mH$ defines an extremal face Σ of dimension at least two, containing $[V]$. By [15, Lemma 7.2] there exists an extremal ray $R \in \Sigma$ whose exceptional locus is contained in the indeterminacy locus B of the $\text{rc}(\mathcal{V})$ -fibration. Since $(K_X + mH) \cdot R = 0$, the length $l(R)$ is greater than or equal to m . Let F be a non-trivial fiber of the contraction associated to R ; then $\dim F \geq m + 1$ by Proposition (1.3), since the contraction associated to R is small, being $\dim B \leq \dim X - 2$.

By part (2) of Lemma (2.4), the dimension of $\text{ChLocus}(\mathcal{V})_F$ is

$$\dim \text{ChLocus}(\mathcal{V})_F \geq \dim F + f_V + 1.$$

As the $\text{rc}(\mathcal{V})$ -equivalence classes are either contained in B or have empty intersection with it, $\text{ChLocus}(\mathcal{V})_F \subset B$. Therefore we get

$$\dim X - 2 \geq \dim B \geq \dim \text{ChLocus}(\mathcal{V})_F \geq f_V + m + 2 \geq \dim X - 1,$$

which is a contradiction. $\square$

As the last preparatory step, we consider the following special case.

Lemma 4.2. *Let V be a quasi-unsplit dominating family of rational curves on a smooth complex projective variety X . Denote by f_V the dimension of a general $\text{rc}(\mathcal{V})$ -equivalence class. Assume that there exists an extremal ray R , independent from $[V]$, whose associated contraction has a fiber F such that $\dim F + f_V \geq \dim X$. Then $\dim F + f_V = \dim X$ and $\text{NE}(X) = \langle [V], R \rangle$. In particular $\rho_X = 2$.*

Proof. By part (1) of Lemma (2.4) we have

$$\dim X \geq \dim \text{ChLocus}(\mathcal{V})_F \geq f_V + \dim F,$$

hence $\dim F + f_V = \dim X$ and $\text{ChLocus}(\mathcal{V})_F = X$; so the assertion follows by Corollary (1.11). $\square$

Theorem 4.3. *Let (X, H) be a polarized manifold with a dominating family of rational curves V such that $H \cdot V = 1$. If $-K_X \cdot V \geq \frac{\dim X - 1}{2}$, then $[V]$ spans an extremal ray of $\text{NE}(X)$.*

Proof. Let B be the indeterminacy locus of the $\text{rc}(\mathcal{V})$ -fibration $q: X \dashrightarrow Y$, let D be a very ample divisor on $q(X \setminus B)$ and let $\widehat{D} := \overline{q^{-1}D}$.

Denote by m the anticanonical degree of V and by f_V the dimension of a general $\text{rc}(\mathcal{V})$ -equivalence class. Note that, since V is a dominating family, we have $m \geq 2$. By Proposition (1.2) $\dim \text{Locus}(V_x) \geq -K_X \cdot V - 1 = m - 1$; since a general fiber of the $\text{rc}(\mathcal{V})$ -fibration contains $\text{Locus}(V_x)$ for every point x in it, we have $f_V \geq m - 1$.

If $K_X + mH$ is nef, then the assertion follows by Proposition (4.1); therefore we can assume that $K_X + mH$ is not nef.

Let R be an extremal ray such that $(K_X + mH) \cdot R < 0$ and let φ_R be the associated contraction. Notice that R has length $l(R) \geq m + 1$, hence every non-trivial fiber of φ_R has dimension $\geq m$ by Proposition (1.3). On the other hand, by Lemma (4.2) we can confine to assume that all fibers of φ_R have dimension $\leq m + 1$. In particular this implies that $H \cdot R = 1$. Indeed, if this were not the case, we would have $l(R) \geq 2m + 1$, hence every non-trivial fiber of φ_R would have dimension $\geq 2m > m + 1$, by Proposition (1.3) and the fact that $m \geq 2$.

If the Picard number of X is one the theorem is clearly true, so we can assume that $\rho_X \geq 2$. Now we split up the proof in two cases, according to the value of ρ_X : first we consider the case $\rho_X = 2$ and then the general one.

Case (a) $\rho_X = 2$.

The proof is based on different arguments, depending on the dimension of the fibers of the contraction associated to the extremal ray R .

Case (a1) The contraction φ_R admits an $(m + 1)$ -dimensional fiber F .

Consider $X_F := \text{ChLocus}(\mathcal{V})_F$. We have, by Corollary (1.11), that $\text{NE}^X(X_F) = \langle [V], R \rangle$ and, by Lemma (2.4), that

$$\dim X_F \geq \dim F + f_V \geq (m + 1) + (m - 1) \geq \dim X - 1.$$

If $X_F = X$, then the statement is proved. So we can assume that an irreducible component $\overline{X}_F$ of X_F is a divisor and thus that $f_V = m - 1$. Notice that $\overline{X}_F \cdot V = 0$, otherwise we would have $X_F = X$; in particular it follows that $\overline{X}_F$ is closed with respect to $\text{rc}(\mathcal{V})$ -equivalence, so $X_F = \overline{X}_F$.

Consider now the intersection number of X_F with curves in R ; since $\rho_X = 2$ and $X_F \cdot V = 0$ we cannot have also $X_F \cdot R = 0$.

Let us show that we cannot have $X_F \cdot R < 0$, too.

Assume by contradiction that this is the case. Then $\text{Exc}(R) \subset X_F$, so φ_R is divisorial by Proposition (1.3). By the same proposition, recalling that we are assuming that all the fibers of φ_R have dimension $\leq m + 1$, every non-trivial fiber has dimension $m + 1$.

Then φ_R is the blow-up of a smooth variety X' along a smooth center T by [2, Theorem 4.1 (iii)]. The dimension of the center is $\dim T = (n - 1) - (m + 1) \leq m - 1 = f_V$. We can apply part (4) of Lemma (3.1) and we get $\rho_X = \rho_{X'} + 1 > 2$, so we reach a contradiction.

Therefore $X_F \cdot R > 0$. It follows that $(X_F)|_{X_F}$ is nef and thus, by Lemma (1.12), X_F is nef.

As $X_F \cdot V = 0$ and $\rho_X = 2$, X_F is the supporting divisor of an elementary contraction of X whose associated extremal ray is spanned by $[V]$.

Case (a2) The contraction φ_R is equidimensional with m -dimensional fibers.

By Proposition (1.3), φ_R is of fiber type and $l(R) = m + 1$. Hence, by [11, Lemma 2.12], X is a projective bundle over a smooth variety Y , *i.e.* $X = \mathbb{P}_Y(\mathcal{E})$, where $\mathcal{E} = (\varphi_R)_* H$.

Notice that Y has Picard number one and is covered by rational curves - the images

of the curves parametrized by V - therefore Y is a Fano manifold.
By the canonical bundle formula for projective bundles we have

$$K_X + (m+1)H = \varphi_R^*(K_Y + \det \mathcal{E}).$$

In particular, if C_V is a curve among those parametrized by V , by the projection formula we can compute

$$(K_Y + \det \mathcal{E}) \cdot (\varphi_R)_*(C_V) = (K_X + (m+1)H) \cdot C_V = 1.$$

It follows that $K_Y + \det \mathcal{E}$ is the ample generator of $\text{Pic}(Y)$ and that $(K_Y + \det \mathcal{E}) \cdot \varphi_R(C_V) = 1$.

The ampleness of $\mathcal{E}$ implies that $\det \mathcal{E} \cdot \varphi_R(C_V) \geq m+1$; therefore $-K_Y \cdot \varphi_R(C_V) \geq m$, hence the index r_Y of Y is greater than or equal to m .

If $r_Y = m$, denoted by l a rational curve of minimal degree in Y , then $\det \mathcal{E} \cdot l = m+1$; moreover, the splitting type of $\mathcal{E}$, which is ample and of rank $m+1$, on rational curves of minimal degree is uniform of type $(1, \dots, 1)$.

We can thus apply [3, Proposition 1.2], so we obtain that $X \simeq \mathbb{P}^m \times Y$. It follows that the curves of V are contained in the fibers of the first projection and that $[V]$ spans an extremal ray.

Therefore we are left with $r_Y \geq m+1$. Recalling that $\dim Y = \dim X - m \leq m+1$, by the Kobayashi–Ochiai Theorem ([13]) we get that Y is a hyperquadric or a projective space.

Assume by contradiction that $[V]$ does not span an extremal ray of X .

By part (3) of Lemma (2.2) there exists a curve $C \subset B$, whose numerical class is not proportional to $[V]$, such that $\widehat{D} \cdot C \leq 0$. Actually, since $\rho_X = 2$ and $\widehat{D} \cdot V = 0$, we have $\widehat{D} \cdot C < 0$.

By part (2) of Lemma (2.4), there exists a component X_C of $\text{ChLocus}(\mathcal{V})_C$ such that $C \subset X_C$ and $\dim X_C \geq f_V + \dim C + 1 \geq m+1$.

By Lemma (1.10) $\widehat{D}$ has non positive intersection number with every curve in X_C and it is trivial only on curves which are numerically proportional to $[V]$.

Since $\widehat{D} \cdot R > 0$, we have that φ_R does not contract curves in X_C , hence $\dim Y \geq \dim X_C \geq m+1$ and so $\dim Y = \dim X_C = m+1$.

Moreover, as $C \subset X_C$ and C is not numerically proportional to $[V]$, we get that X_C is not $\text{rc}(\mathcal{V})$ -connected; in particular, for every point c of C , the intersection X_c of the $\text{rc}(\mathcal{V})$ -equivalence class containing c with X_C has dimension $\leq m$.

We claim that there exists a line l in Y which is not contained in the image of $\varphi_R(X_c)$ for any $c \in C$. Notice that, since φ_R does not contract curves in X_C , through a general point y in Y there is a finite number of such subvarieties.

If $Y \simeq \mathbb{P}^{m+1}$, a line joining y with a point outside the union of these subvarieties has the required property.

If $Y \simeq \mathbb{Q}^{m+1}$, then for a general y the locus of the lines through y is not contained in $\varphi_R(X_c)$ for any $c \in C$, hence a general line through y has the required property.

The splitting type of $\mathcal{E}$ on this line is one of the following: $(2, 1, \dots, 1)$ if $Y \simeq \mathbb{Q}^{m+1}$ and either $(3, 1, \dots, 1)$ or $(2, 2, 1, \dots, 1)$ if $Y \simeq \mathbb{P}^{m+1}$. Recalling that $m \geq 2$ we have that, among the summands of $\mathcal{E}_l$ there is at least one $\mathcal{O}_{\mathbb{P}^1}(1)$.

Consider $\mathbb{P}_l(\mathcal{E}|_l)$; its cone of curves is generated by the class of a line in a fiber of the projection onto l and the class of a minimal section C_0 . By the discussion above we have that $H \cdot C_0 = 1$. Moreover, $\varphi_R^*(K_Y + \det \mathcal{E}) \cdot C_0 = 1$, hence $[C_0] = [V]$; in

particular $\widehat{D}$ is nef on $\mathbb{P}_l(\mathcal{E}|_l)$.

Consider an irreducible curve in $\mathbb{P}_l(\mathcal{E}|_l) \cap X_C$; by our choice of l , this curve is not contained in a $\text{rc}(\mathcal{V})$ -equivalence class contained in X_C , so it is negative with respect to $\widehat{D}$, a contradiction. The case $\rho_X = 2$ is thus completed.

Case (b) $\rho_X > 2$.

Notice that, in view of Corollary (2.3), we can confine to assume that $B \neq \emptyset$; moreover, by part (3) of Lemma (2.2), we can also assume the existence of a curve $C \subset B$ such that $[C]$ is not proportional to $[V]$ and $\widehat{D} \cdot C \leq 0$.

We claim that $K_X + (m+1)H$ is nef.

Assume by contradiction that $K_X + (m+1)H$ is not nef. Let τ be a ray such that $(K_X + (m+1)H) \cdot \tau < 0$, denote by Γ_τ a rational curve of minimal anticanonical degree in τ and by φ_τ the contraction associated to τ . Notice that τ has length $l(\tau) \geq m+2$, hence every non-trivial fiber of φ_τ has dimension $\geq m+1$ by Proposition (1.3). Moreover, φ_τ cannot have fibers of dimension $> m+1$, otherwise, by Lemma (4.2), we would have $\rho_X = 2$.

Therefore every non-trivial fiber of φ_τ has dimension $m+1$. In view of Proposition (1.3), we thus get that φ_τ is of fiber type and that the length of τ is $l(\tau) = m+2$; this last fact gives $H \cdot \Gamma_\tau = 1$.

Let W_τ be a family of deformations of Γ_τ .

Since B is not empty, there are $\text{rc}(\mathcal{V})$ -equivalence classes of dimension $\geq f_V + 1 \geq m$; let G be one of these classes.

By part (2) of Lemma (2.4) we have

$$\dim \text{ChLocus}(\mathcal{W}_\tau)_G \geq \dim G + (m+1) = 2m+1 \geq \dim X,$$

so by Lemma (1.9) we deduce $\rho_X = 2$, a contradiction which proves the nefness of $K_X + (m+1)H$.

Notice that $K_X + (m+1)H$ is not ample, as $(K_X + (m+1)H) \cdot R = 0$. Let Σ be the extremal face contracted by $K_X + (m+1)H$. We now consider separately two cases, depending on the existence in Σ of a fiber type extremal ray.

Case (b1) There exists a fiber type extremal ray ϑ in Σ .

Let φ_ϑ be the contraction associated with ϑ and denote by W_ϑ a (dominating) family of deformations of a rational curve of minimal anticanonical degree in ϑ .

By part (2) of Lemma (2.4), there exists a component X_C of $\text{ChLocus}(\mathcal{V})_C$ such that $C \subset X_C$ and $\dim X_C \geq f_V + 2$.

According to Lemma (1.10), every curve in X_C can be written as $\alpha[C] + \beta[V]$ with $\alpha \geq 0$; in particular, since $\widehat{D} \cdot V = 0$ by Lemma (2.2), it follows that $\widehat{D}$ is not positive on any curve contained in X_C . By the same lemma $\widehat{D} \cdot W_\vartheta > 0$, hence $[W_\vartheta] \notin \text{NE}^X(X_C)$. Therefore part (1) of Lemma (2.4) gives

$$\dim \text{ChLocus}(\mathcal{W}_\vartheta)_{X_C} \geq \dim X_C + f_{W_\vartheta} \geq f_V + 2 + m \geq \dim X,$$

where f_{W_ϑ} is the dimension of the general $\text{rc}(\mathcal{W}_\vartheta)$ -equivalence class.

Moreover, by Lemma (1.10) the class of every curve in X can be written as

$$\alpha[C] + \beta[V] + \gamma[W_\vartheta]$$

with $\alpha \geq 0$.

In particular, since we are assuming $\rho_X > 2$, this implies that every curve $C \subset B$ not numerically proportional to $[V]$ such that $\widehat{D} \cdot C \leq 0$ cannot be contained in the

plane Π spanned by $[V]$ and $[W_\vartheta]$.

It follows that $\widehat{D}$ is nef on the two-dimensional face $\Pi \cap \text{NE}(X)$, as $\widehat{D} \cdot \gamma < 0$ implies $\gamma \in B$. Recalling that $\widehat{D} \cdot W_\vartheta > 0$, we thus have that $[V]$ spans an extremal ray.

Case (b2) Every ray in Σ is birational.

Let η be any ray in Σ . By Proposition (1.3), for every non-trivial fiber of its associated contraction φ_η we have $\dim F \geq l(\eta) \geq m+1$. Recalling that at the beginning of the proof we observed that we can assume $\dim F \leq m+1$, we have $\dim F = m+1 = l(\eta)$. Therefore, by Proposition (1.3), φ_η is a divisorial contraction, hence, by [2, Theorem 4.1 (iii)], is the blow-up of a smooth variety along a smooth center T of dimension $(n-1) - (m+1) \leq m-1$.

Let E be the exceptional divisor of φ_η . We claim that $E \cdot V = 0$.

By part (2) of Lemma (2.4), there exists a component X_C of $\text{ChLocus}(\mathcal{V})_C$ such that $C \subset X_C$ and $\dim X_C \geq f_V + 2$.

By Lemma (1.10) $\widehat{D}$ has non positive intersection number with every curve in X_C . If $E \cap X_C \neq \emptyset$, then there is a fiber F of φ_η meeting X_C . Counting dimensions, we find that $\dim(F \cap X_C) \geq 1$, which is a contradiction as $\widehat{D} \cdot \eta > 0$. So $E \cap X_C = \emptyset$, whence $E \cdot V = 0$, proving the claim.

Therefore E contains $\text{rc}(\mathcal{V})$ -equivalence classes and $\dim T \geq f_V$, since φ_η is finite to one on $\text{rc}(\mathcal{V})$ -equivalence classes. Recalling that $f_V \geq m-1$ we derive $\dim T = f_V = m-1$.

Assume that $\dim \Sigma \geq 2$ and let E_1, E_2 be the exceptional loci of two different extremal rays η_1, η_2 in Σ ; we claim that $E_1 \cap E_2 = \emptyset$.

To prove the claim, notice that $E_i = \text{ChLocus}(\mathcal{V})_{F_i}$ for any fiber F_i of η_i , since E_i is closed with respect to $\text{rc}(\mathcal{V})$ -equivalence and, by part (1) of Lemma (2.4), we can compute

$$\dim \text{ChLocus}(\mathcal{V})_{F_i} \geq m+1 + f_V = \dim E_i.$$

So, Corollary (1.11) gives $\text{NE}(E_i) = \langle [V], \eta_i \rangle$, hence E_1 cannot contain a curve in a fiber of the contraction associated to η_2 ; since the dimension of the fibers is ≥ 2 this implies that $E_1 \cdot R_2 = 0$. But then, if $E_1 \cap E_2 \neq \emptyset$, then E_1 would contain the fibers of φ_2 which it meets. So we reach a contradiction, proving the claim.

Therefore the contraction $\sigma: X \rightarrow X'$ of the face Σ verifies the assumptions of Lemma (3.1), hence there exist an ample line bundle H' on X' and an unsplit dominating family V' on X' such that $H' \cdot V' = 1$ and $-K_{X'} \cdot V' = -K_X \cdot V \geq \frac{\dim X' - 1}{2}$.

Denote by $f_{V'}$ the dimension of the general $\text{rc}(\mathcal{V}')$ -equivalence class. Since a general fiber of the $\text{rc}(\mathcal{V}')$ -fibration contains $\text{Locus}(V'_{x'})$, we have $f_{V'} \geq \dim \text{Locus}(V'_{x'}) - 1 \geq m-1$.

Consider the adjoint divisor $K_{X'} + mH'$; if it is nef, or an extremal ray R' such that $(K_{X'} + mH') \cdot R' < 0$ has a fiber of dimension $\geq m+2$, then $[V']$ spans an extremal ray by Proposition (4.1) or by Lemma (4.2), so $[V]$ spans an extremal ray by Lemma (3.1).

Let us show that the remaining case does not happen.

Assume that there is an extremal ray R' such that $(K_{X'} + mH') \cdot R' < 0$ and every fiber of the associated contraction has dimension $\leq m+1$. In particular we have $H' \cdot R' = 1$, otherwise we would have $l(R') \geq 2m+1$, hence every non-trivial fiber of the associated contraction would have dimension $\geq 2m > m+1$ by Proposition

(1.3). Moreover, we have $(K_{X'} + (m+1)H') \cdot R' \leq 0$, since $l(R') \geq m+1$. On the other hand, recalling that $\sigma^*H' = H - \sum E_i$ and that $\sigma^*K_{X'} = K_X + \sum (m+1)E_i$, we have

$$\sigma^*(K_{X'} + (m+1)H') = K_X + (m+1)H,$$

so, by the projection formula, $K_{X'} + (m+1)H'$ is ample on X' , a contradiction. $\square$

Corollary 4.4. *Let (X, H) be a polarized manifold of dimension at most five, with a dominating family of rational curves V such that $H \cdot V = 1$. Then $[V]$ spans an extremal ray of $\text{NE}(X)$.*

5. AN EXAMPLE

In the paper [5], an application of the results about extremality of families of lines was a relative version of a theorem proved in [17], which was the first step towards a conjecture of Mukai for Fano manifolds.

This conjecture states that, for a Fano manifold X , denoted by ρ_X its Picard number and by r_X its index, we have

$$\rho_X(r_X - 1) \leq \dim X.$$

In [17, Theorem B] it was proved that, if $r_X \geq \frac{\dim X}{2} + 1$, then $\rho_X = 1$ unless $X \simeq \mathbb{P}^{\dim X/2} \times \mathbb{P}^{\dim X/2}$.

In [5, Theorem 3.1.1] it was proved that a fiber type contraction $\varphi: X \rightarrow Y$ supported by $K_X + mL$ with $m \geq \frac{\dim X}{2} + 1$ is elementary, unless $X \simeq \mathbb{P}^{\dim X/2} \times \mathbb{P}^{\dim X/2}$.

In the last few years some progress has been made towards Mukai conjecture; in particular it was proved in [1] that it holds for a Fano manifold with (pseudo)index greater than or equal to $\frac{\dim X}{3} + 1$ if X admits an unsplit dominating family of rational curves.

It is therefore natural to ask if the corresponding relative statement is true, namely, given a fiber type contraction $\varphi: X \rightarrow Y$, corresponding to an extremal face Σ , supported by $K_X + mL$ with $m \geq \frac{\dim X}{3} + 1$ and such that there is an unsplit dominating family V whose curves are contracted by φ , is there a bound on the dimension of Σ ?

The answer to this question is negative, as we will show with an example in which $m = \frac{\dim X}{2}$; it follows that [5, Theorem 3.1.1] cannot be improved.

Example 5.1. *Let Z be a smooth variety of dimension $k+2$ and denote by Y the product $Z \times \mathbb{P}^k$; denote by p_1 and p_2 the projections onto the factors. Let $\{z_i\}_{i=1, \dots, t}$ be points of Z and denote by F_i the fibers of p_1 over z_i . Let $\sigma: X \rightarrow Y$ be the blow-up of Y along the union of F_i . The canonical bundle of X is*

$$(5.1.1) \quad K_X = \sigma^*K_Y + (k+1) \sum_i E_i = \sigma^*(p_1^*K_Z + p_2^*K_{\mathbb{P}^k}) + (k+1) \sum_i E_i;$$

denoting by $\mathcal{H} := (p_2 \circ \sigma)^*\mathcal{O}_{\mathbb{P}^k}(1)$ and by $L' := \mathcal{H} - \sum E_i$, we can rewrite formula (5.1.1) as

$$K_X + (k+1)L' = \sigma^*(p_1^*K_Z)$$

It is easy to check that L' is $(p_1 \circ \sigma)$ -ample. Let $A \in \text{Pic}(Z)$ be an ample line bundle; then $L := L' + \sigma^*(p_1^*A)$ is an ample line bundle on X ; moreover $L \cdot l = 1$ for a line l in the strict transform of a fiber F of p_1 not contained in the center of σ .

The contraction $p_1 \circ \sigma$ is supported by $K_X + (k+1)L = K_X + \frac{\dim X}{2}L$ and contracts a face of dimension $t+1$.

Remark 5.2. The difference between the relative and the absolute case is given by the existence of minimal horizontal dominating families of rational curves for proper morphisms defined on a open subset of a Fano manifold (for the definition and the references see [1, Remark 6.4]). Such families do not exist in general in the relative case.

ACKNOWLEDGEMENTS

We learned of the results about extremality of families of lines in [5] from an interesting series of lectures given by Paltin Ionescu.

REFERENCES

- [1] Marco Andreatta, Elena Chierici, and Gianluca Occhetta. Generalized Mukai conjecture for special Fano varieties. *Cent. Eur. J. Math.*, 2(2):272–293, 2004.
- [2] Marco Andreatta and Jarosław A. Wiśniewski. A note on nonvanishing and applications. *Duke Math. J.*, 72(3):739–755, 1993.
- [3] Marco Andreatta and Jarosław A. Wiśniewski. On manifolds whose tangent bundle contains an ample subbundle. *Invent. Math.*, 146(1):209–217, 2001.
- [4] Mauro C. Beltrametti and Paltin Ionescu. On manifolds swept out by high dimensional quadrics. *Math. Z.*, 2008. to appear.
- [5] Mauro C. Beltrametti, Andrew J. Sommese, and Jarosław A. Wiśniewski. Results on varieties with many lines and their applications to adjunction theory. In *Complex algebraic varieties (Bayreuth, 1990)*, volume 1507 of *Lecture Notes in Math.*, pages 16–38. Springer, Berlin, 1992.
- [6] Laurent Bonavero, Cinzia Casagrande, and Stéphane Druel. On covering and quasi-unsplit families of rational curves. *J. Eur. Math. Soc.*, 9(1):45–57, 2007.
- [7] F. Campana. Coréduction algébrique d’un espace analytique faiblement kählerien compact. *Invent. Math.*, 63(2):187–223, 1981.
- [8] F. Campana. Connexité rationnelle des variétés de Fano. *Ann. Sci. École Norm. Sup. (4)*, 25(5):539–545, 1992.
- [9] Elena Chierici and Gianluca Occhetta. The cone of curves of Fano varieties of coindex four. *Internat. J. Math.*, 17(10):1195–1221, 2006.
- [10] Olivier Debarre. *Higher-dimensional algebraic geometry*. Universitext. Springer-Verlag, New York, 2001.
- [11] Takao Fujita. On polarized manifolds whose adjoint bundles are not semipositive. In *Algebraic geometry, Sendai, 1985*, volume 10 of *Adv. Stud. Pure Math.*, pages 167–178. North-Holland, Amsterdam, 1987.
- [12] Paltin Ionescu. Generalized adjunction and applications. *Math. Proc. Cambridge Philos. Soc.*, 99(3):457–472, 1986.
- [13] Shoshichi Kobayashi and Takushiro Ochiai. Characterizations of complex projective spaces and hyperquadrics. *J. Math. Kyoto Univ.*, 13:31–47, 1973.
- [14] János Kollár. *Rational curves on algebraic varieties*, volume 32 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, Berlin, 1996.
- [15] Carla Novelli and Gianluca Occhetta. Projective manifolds containing a large linear subspace with nef normal bundle. Preprint math.AG 0712.3406, 2007.
- [16] Gianluca Occhetta. A characterization of products of projective spaces. *Canad. Math. Bull.*, 49:270–280, 2006.
- [17] Jarosław A. Wiśniewski. On a conjecture of Mukai. *Manuscripta Math.*, 68(2):135–141, 1990.
- [18] Jarosław A. Wiśniewski. On contractions of extremal rays of Fano manifolds. *J. Reine Angew. Math.*, 417:141–157, 1991.

DIPARTIMENTO DI MATEMATICA “F. CASORATI”,
UNIVERSITÀ DI PAVIA,
VIA FERRATA 1,
I-27100 PAVIA

E-mail address: `carla.novelli@unipv.it`

DIPARTIMENTO DI MATEMATICA,
UNIVERSITÀ DI TRENTO,
VIA SOMMARIVE 14,
I-38050 Povo (TN)

E-mail address: `gianluca.occhetta@unitn.it`